

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
3 January 2002 (03.01.2002)

PCT

(10) International Publication Number
WO 02/00875 A2

Document AP9
Appl. No. 09/517,466

(51) International Patent Classification⁷: C12N 15/10

(74) Agent: CHEN, Shirley; Wilson Sonsini Goodrich & Rosati, 650 Page Mill Road, Palo Alto, CA 94304-1050 (US).

(21) International Application Number: PCT/US01/19770

(22) International Filing Date: 20 June 2001 (20.06.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
09/606,323 28 June 2000 (28.06.2000) US

(63) Related by continuation (CON) or continuation-in-part (CIP) to earlier application:
US 09/606,323 (CON)
Filed on 28 June 2000 (28.06.2000)

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (*for all designated States except US*): PRO-TEMATION, INC. [US/US]; 520 Breck Court, Benicia, CA 94510 (US).

Published:

— without international search report and to be republished upon receipt of that report

(72) Inventor; and

(75) Inventor/Applicant (*for US only*): CLARK, Robin [US/US]; 520 Breck Court, Benicia, CA 94510 (US).

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: COMPOSITIONS AND METHODS FOR GENERATING EXPRESSION VECTORS THROUGH SITE-SPECIFIC RECOMBINATION

(57) Abstract: Compositions, kits, and methods are provided for use in a recombinational cloning or subcloning methods for constructing expression vectors which comprise: ligating a library of double-stranded linear donor DNAs, where each member of the library includes a donor DNA sequence, with a double-stranded linear driver DNA which includes a promoter sequence and a donor recombination site to form a single circular donor DNA, the single circular donor DNA not including an origin of replication, where the donor DNA sequence is under the transcriptional control of the promoter; and contacting the circular donor DNA and a circular acceptor vector in the presence of a recombinase to form a single fused circular vector, the circular acceptor vector comprising an origin of replication and an acceptor recombination site capable of recombining with the circular donor DNA.

WO 02/00875 A2

BEST AVAILABLE COPY

COMPOSITIONS AND METHODS FOR GENERATING EXPRESSION VECTORS THROUGH SITE-SPECIFIC RECOMBINATION

5

Field of the Invention

This invention relates to recombinant DNA technology, nucleic acids, vectors and methods for use in a recombinational cloning or subcloning, and more specifically for constructing expression vectors by using recombination proteins in vitro or in vivo through site-specific recombination.

10

Description of Related Art

Recombinant DNA technology, also called gene cloning or molecular cloning, is widely used to transfer genetic information, i.e. DNA, from one organism to another. A typical recombinant DNA experiment often follows the following procedure. First, the DNA (e.g., the cloned DNA, insert DNA, target DNA, or foreign DNA) from a donor organism is extracted, enzymatically cleaved (or cut/digested), and joined (ligated) to another DNA entity (e.g. a cloning vector) to form a new, recombinant DNA molecule (or cloning vector-insert DNA construct). Second, this cloning vector-insert DNA construct is transferred into and maintained within a host cell, such as transformation of a bacterial host cell by the construct. Third, those host cells that take up the DNA construct (transformed cells) are identified and selected from those that do not. In addition, if required, a DNA construct can be prepared to ensure that the protein product that is encoded by the cloned DNA sequence is produced by the host cell.

15

20

25

30

Accordingly, this traditional cloning methods using restriction enzymes and ligase can be time consuming, especially when a specific

expression vector is required for transferring the target gene into a heterologous host cell, such as a mammalian cell. The specific expression vector may not contain matching restriction sites for the donor DNA. Extensive reengineering of the expression vector may be required to introduce the matching restriction sites into the vector so that the vector and the insert DNA can be ligated to produce the final construct. Alternatively, multiple restriction enzymes may have to be employed to generate an insert DNA having suitable restriction sites for ligation with the vector. In this case, reaction conditions for each restriction enzyme may differ such that it is often necessary to perform a few separate restriction digestion reactions to obtain the desired insert. Further, the efficiency of direct ligation between the vector and insert may be very low, especially between large fragments. As a result, the whole procedure is tedious, and the final yield of the correctly ligated construct can be low.

Site-specific recombination represents another useful method of recombinant DNA technology. This method employs a site-specific recombinase, an enzyme which catalyzes the exchange of DNA segments at specific recombination sites. Site-specific recombinases present in some viruses and bacteria, and have been characterized to have both endonuclease and ligase properties. These recombinases, along with associated proteins in some cases, recognize specific sequences of bases in DNA and exchange the DNA segments flanking those segments. Landy, A. (1993) *Current Opinion in Biotechnology* 3:699-707.

A typical site-specific recombinase is Cre recombinase. Cre is a 38-kDa product of the *cre* (cyclization recombination) gene of bacteriophage P1 and is a site-specific DNA recombinase of the Int family. Sternberg, N. et al. (1986) *J. Mol. Biol.* 187: 197-212. Cre

recognizes a 34-bp site on the P1 genome called loxP (locus of X-over of P1) and efficiently catalyzes reciprocal conservative DNA recombination between pairs of loxP sites. The loxP site consists of two 13-bp inverted repeats flanking an 8-bp nonpalindromic core region.

5 Cre-mediated recombination between two directly repeated loxP sites results in excision of DNA between them as a covalently closed circle. Cre-mediated recombination between pairs of loxP sites in inverted orientation will result in inversion of the intervening DNA rather than excision. Breaking and joining of DNA is confined to discrete positions
10 within the core region and proceeds on strand at a time by way of transient phosphotyrosine DNA-protein linkage with the enzyme. Other examples of site-specific recombination systems include the integrase/att system from bacteriophage λ and the FLP/FRT system from the *Saccharomyces cerevisiae* 2 μ i circle plasmid.

15

These site-specific recombination systems have been used in vivo to facilitate recombination between different vectors. Waterhouse et al. used an in vivo method to join light and heavy chains of an antibody. The light and heavy chains were cloned in different phage
20 vectors between loxP and loxP 511 sites that were used to transform new *E. coli* cells. Waterhouse, P. et al. (1993) Nucleic Acid Res. 21:2265-2266. Cre acted on two parental molecules, one plasmid and another phage, in the host cells to produce four products in equilibrium: two different cointegrates (produced by recombination at either loxP or
25 loxP511 sites), and two daughter molecules, one of which was the desired product. Schlake and Bode used an *in vivo* method to exchange expression cassettes at defined chromosomal locations, each flanked by a wild type and spacer-mutated FRT recombination site. Schlake and Bode (1994) Biochemistry 33:12746-12751. A double-
30 reciprocal crossover was mediated in cultured mammalian cells by using the FLP/FRT system for site-specific recombination. Aoki et al. used a

shuttle plasmid (pAdMCS) that carried a gene of interest, a loxP site, the adenoviral 5-LTR and packaging signal 0 to 1 mu, and a multiple cloning site. Aoki et al. (1999) Mol. Med. 5:224-231. The shuttle plasmid was linearized by a restriction enzyme NheI and recombined with ClaI-digested adenoviral cosmid *in vitro*. Cre recombinase produced the full-length recombinant adenoviral vector *in vitro* by an exchange of region distal to the loxP site linearized in these two molecules.

10 **SUMMARY OF THE INVENTION**

The present invention relates to compositions, kits, and methods for use in a recombinational cloning or subcloning. In particular, the present invention provides novel methods for constructing expression vectors by using site-specific recombinases *in vitro*. These method may be used for high throughput screening of genes, functional genomics and other human genome projects.

In one aspect, the present invention provides a double-stranded circular donor DNA for transferring a donor DNA sequence into expression vectors. The circular donor DNA comprises: a donor DNA sequence; a donor recombination site; at least one selectable marker, the circular donor DNA not including an origin of replication.

The donor DNA sequence may be any gene of interest or any synthetic DNA sequence which is needed to be transferred into an expression vector. For example the donor DNA segment may be a sequence derived from cDNA of a particular gene or one of the members of a cDNA library. The donor DNA may also be a genomic DNA that contains the coding region interrupted with non-coding sequences.

In another aspect, the present invention also provides a library of double-stranded circular donor DNAs that may be used for high throughput screening. The library of double-stranded circular DNA
5 comprises: a donor DNA sequence which varies within a library of donor DNA sequences; a donor recombination site; and at least one selectable marker, the circular donor DNA not including an origin of replication.

The library of donor DNA sequences may be a library of cDNA or
10 genomic DNA derived from any desirable sources. For example, the library of donor DNA sequences may be a cDNA library from single human chromosomes.

The circular donor DNA may further comprise a promoter
15 sequence that controls expression of the donor DNA sequence. The promoter may be any array of DNA sequences that interact specifically with cellular transcription factors to regulate transcription of the downstream gene. The promoter may be derived from any organism, such as bacteria, yeast, insect and mammalian cells and viruses.
20 Examples of the promoter include, but are not limited to, *E. coli lac* and *trp* operons, the *tac* promoter, the bacteriophage λ *p^L* promoter, bacteriophage T7 and SP6 promoters, β -actin promoter, insulin promoter, human cytomegalovirus (CMV) promoter, HIV-LTR (HIV-long terminal repeat), Rous sarcoma virus RSV-LTR, simian virus SV40
25 promoter, baculoviral polyhedrin and p10 promoter.

The promoter may also be an inducible promoter that regulates the expression of downstream gene in a controlled manner. Examples of inducible promoters include, but are not limited to, the bacterial dual
30 promoter (activator/repressor expression system) which regulates gene expression in mammalian cells under the control of tetracycline and its

analogs and promoters that regulate gene expression under the control of factors such as heat shocks, steroid hormones, heavy metals, phorbol ester, the adenovirus E1A element, interferon, or serum.

5 The donor recombination site may be any segment or arrays of DNA sequence recognized by a site-specific recombinase which catalyzes site-specific fusion between the circular donor DNA and an acceptor vector. The site-specific recombinase may be a recombinase, a transposase or an integrases.

10

 In one variation, the recombination site is a lox site that is recognized by the Cre recombinase of bacteriophage P1. Example of lox site includes, but are not limited to, loxB, loxL, loxR, loxP [SEQ ID NO:1], loxP3, loxP23, lox Δ 86, lox Δ 117, loxP511 [SEQ ID NO:2], and
15 loxC2 [SEQ ID NO:3].

 In another variation, the recombination site is a recombination site that is recognized by a recombinases other than Cre. Examples of the non-Cre recombinases include, but are not limited to, site-specific
20 recombinases include: att sites recognized by the Int recombinase of bacteriophage λ (e.g. att1, att2, att3, attP, attB, attL, and attR), the FRT sites recognized by FLP recombinase of the 2pi plasmid of *Saccharomyces cerevisiae*, the recombination sites recognized by the resolvase family, and the recombination site recognized by transposase
25 of *Bacillus thuringiensis*.

 The example of site-specific recombinase include, but are not limited to, bacteriophage P1 Cre recombinase, yeast FLP recombinase, Inti integrase, bacteriophage λ , phi 80, P22, P2, 186, and P4
30 recombinase, Tn3 resolvase, the Hin recombinase, and the Cin recombinase, *E. coli* xerC and xerD recombinases, *Bacillus thuringiensis*

recombinase, *TpnI* and the β -lactamase transposons, and the immunoglobulin recombinases,

5 The selectable marker of the circular donor DNA may be any functional element for facilitating subsequent identification and selection of clones of the recombination product under suitable conditions. The selectable marker may encode any functional element, such as protein, peptide, RNA, binding site for RNA and proteins, or products that provide resistance to organic or inorganic agents. Examples of
10 selectable markers include, but are not limited to, reporter genes such as β -galactosidase (GAL), fluorescent proteins (e.g., GFP, GFP-UV, EFPF, BFP, EBFP, ECFP, EYFP), secreted form of human placental alkaline phosphatase (SEAP), β -glucuronidase (GUS)); resistance genes against antibiotics (e.g. neomycin (G418) or hygromycin resistant
15 gene, puromycin resistant gene), yeast selectable markers *leu2-d* and URA3, apoptosis resistant genes (e.g. baculoviral p35 gene), and antisense oligonucleotides.

20 The circular donor DNA may optionally include an affinity tag for selection and isolation of protein product encoded by the donor DNA segment. Examples of such an affinity tag include, but are not limited to; a polyhistidine tract, polyarginine, glutathione-S-transferase (GST), maltose binding protein (MBP), a portion of staphylococcal protein A (SPA), and various immunoaffinity tags (e.g. protein A) and epitope tags
25 such as those recognized by the EE (Glu-Glu) antipeptide antibodies. The affinity tag may be positioned at either the amino- or carboxy-terminus of the donor DNA.

30 The present invention also provides a circular acceptor vector for generating recombinant expression vector. The vector comprises an origin of replication; and an acceptor recombination site capable of

recombining with a donor DNA. Optionally, the acceptor vector may not include a promoter for regulating expression of the donor DNA.

5 The circular acceptor vector may be any vector that can transform, transfect or transduce a host cell. The acceptor vector may be a plasmid, a phage or a viral vector as long as it is able to replicate in vitro or in a host cell, or to convey the donor DNA to a desired location within a host cell. Examples of host cells include, but are not limited to, bacterial (e.g. *E. coli*, *Bacillus subtilis*, etc.), yeast, animal, plant, and
10 insect cells.

 In one variation, the circular acceptor vector may be a prokaryotic plasmid. Optionally, the acceptor vector may comprise a prokaryotic termination sequence. Examples of the prokaryotic termination
15 sequence include, but are not limited to, the T7 termination sequence, the T_{INT} , T_{LI} , T_{L2} , T_{L3} , TR_1 , TR_2 , T_{6S} termination signals derived from the bacteriophage λ .

 In another variation, the circular acceptor vector may be a
20 mammalian expression vector. The mammalian expression vector contains one or more eukaryotic marker genes, appropriate eukaryotic transcriptional and translational termination signals and a sequence that signals polyadenylation of the transcript messenger RNA (mRNA), and an origin of replication that functions in a mammalian host cell.
25 Examples of the eukaryotic polyadenylation sequence include, but are not limited to, the Herpes simplex virus thymidine kinase polyadenylation sequence, the bovine growth hormone polyadenylation sequence, and the simian virus 40 polyadenylation sequence.

30 Optionally, the eukaryotic expression vector may also carry an origin of replication and selectable marker genes that function in.

bacterial cells, forming a shuttle vector.

5 In yet another variation the circular acceptor includes a promoter for regulating expression of the donor DNA sequence carried by a circular donor DNA of the invention. According to this variation, the recombination site may be placed downstream of the promoter and the transcription initiation site in the acceptor vector.

10 In yet another variation, the circular acceptor may be a yeast expression vector such as a *S. cerevisiae* expression vector. Various types of *S. cerevisiae* expression vector include, but are not limited to, episomal or plasmid vector, integrating vectors, and yeast chromosomes (YACs).

15 In yet another variation, the circular acceptor vector may be a baculovirus DNA, such as wild type or mutant genomes of *Autographa californica multiple nuclear polyhedrosis virus* (AcMNPV) virus.

20 Optionally, a baculoviral acceptor vector according to the present invention may not contain a polyhedrin promoter. Instead, the polyhedrin or the baculoviral p10 promoter can be positioned upstream of the donor DNA sequence of the circular donor DNA of the present invention.

25 The present invention also provides kits for generating recombinant vectors. In one embodiment, the kit comprises: a double-stranded circular donor DNA comprising a donor DNA sequence, a donor recombination site, and at least one selectable marker, the circular donor DNA not including an origin of replication; and a circular
30 acceptor vector comprising an origin of replication and an acceptor recombination site capable of recombining with the circular donor DNA.

In another embodiment, the kit comprises: a library of double-stranded circular donor DNA comprising a donor DNA sequence which varies within a library of donor DNA sequences, a donor recombination site, and at least one selectable marker, the circular donor DNA not including an origin of replication; and a circular acceptor vector comprising an origin of replication and an acceptor recombination site capable of recombining with the circular donor DNA.

In yet another embodiment, the kit comprises: one or more linear donor DNA comprising a donor DNA sequence; a linear driver DNA comprising a promoter sequence, a recombination site, and at least one selectable marker, ligation of the linear donor DNA and the linear driver DNA resulting in a circular donor DNA; and a circular acceptor vector comprising an origin of replication and an acceptor recombination site capable of recombining with the circular donor DNA.

The present invention also provides a method for generating recombinant expression vector *in vitro* through site-specific recombination between a circular donor DNA and circular acceptor DNA, each containing recombination site recognized by the recombinase. The method comprises: contacting a circular double-stranded donor DNA and a circular acceptor vector in the presence of a recombinase under conditions suitable for the circular double-stranded donor DNA and circular acceptor vector to recombine to form a single fused circular vector. In this method, the circular double-stranded donor DNA comprises a donor DNA sequence, a donor recombination site, and at least one selectable marker, but not including an origin of replication. The circular acceptor vector comprises an origin of replication and an acceptor recombination site capable of recombining with the circular donor DNA. The promoter for regulating expression of

the donor DNA may be contained in either the donor DNA or acceptor vector.

5 According to this method, the circular donor DNA containing a site-specific recombination site may be recombined with a circular acceptor vector in the presence of Cre recombinase. The recombination sites on the circular donor DNA and the circular acceptor vector may each contain a lox site.

10 The method may further include steps of transforming, transfecting or transducing a host cell and selecting the correctly fused recombinant vector based on the selectable phenotype conferred by the selectable marker gene on the recombinant vector.

15 The present invention also provides a method for generating recombinant expression vectors from linear DNA segments in vitro. The method comprises: ligating one or more double-stranded linear donor DNA which includes a donor DNA sequence with a double-stranded linear driver DNA which includes a promoter sequence and a donor recombination site to form a single circular donor DNA, the singular
20 circular donor DNA not including an origin of replication, where the donor DNA sequence is under the transcriptional control of the promoter; and contacting the circular donor DNA and a circular acceptor acceptor vector in the presence of a recombinase to form a single fused
25 circular vector. In this method, the circular acceptor vector comprises an origin of replication and an acceptor recombination site capable of recombining with the circular donor DNA.

30 According to this method, the linear donor DNA and linear driver DNA may contain matching restriction sites or other type of annealing sites so as to be ligated to form a circulaized DNA. The linear donor

and driver DNAs may be derived from PCR amplification products.

The present invention also provides a method for high throughput production of recombinant expression vectors from linear DNA segments in vitro. The method comprises: ligating a library of double-stranded linear donor DNAs, where each member of the library includes a donor DNA sequence, with a double-stranded linear driver DNA which includes a promoter sequence and a donor recombination site to form a single circular donor DNA, the singular circular donor DNA not including an origin of replication, where the donor DNA sequence is under the transcriptional control of the promoter; and contacting the circular donor DNA and a circular acceptor vector in the presence of a recombinase to form a single fused circular vector. In this method, the circular acceptor vector comprises an origin of replication and an acceptor recombination site capable of recombining with the circular donor DNA.

According to this method, the library of double-stranded linear donor DNAs may be DNAs amplified from a library of cDNA clones. The library of cDNA clones may be arrayed in a multi-well plate such as 96- and 384-well plates. The library of cDNA clones may be a cosmid or phage library.

Also according to the method, ligating the library of double-stranded linear donor DNAs with a double-stranded linear driver DNA may be performed by Ligation Independent Cloning (LIC). Alternatively, ligating the library of double-stranded linear donor DNAs with a double-stranded linear driver DNA may be performed in the presence of T4 DNA ligase.

The method may further include a step of transferring the

recombinant expression vector into a host and isolating the protein expressed from the vector by affinity tagging. The affinity tagging may be based on a polyhistidine tag (e.g. His₆), a protein tag (e.g., GST, maltose binding protein) or an epitope tag (e.g. an EE ag).

5

The methods of the present invention allow rapid and efficient generation of expression vectors containing the gene of interest without bacterial cloning. Direct ligation of linear donor DNA and linear driver DNA to generate a circular donor DNA allows for efficient cloning of donor DNA such as a cDNA library into an expression vector in an automated and high throughput manner. The methods can be used in a wide variety of high throughput arrays for functional genomics, protein genomics (proteomics), and other human genome projects.

10

15 **BRIEF DESCRIPTION OF FIGURES**

Figure 1 illustrates a process of constructing a library of baculoviral expression vectors through Cre-mediated site-specific recombination.

20

Figure 2 illustrates a process of constructing a baculoviral expression vector for the GUS gene through Cre-mediated site-specific recombination.

25

DETAILED DESCRIPTION OF THE INVENTION

30

The present invention provides reagents, kits and methods for use in a recombinational cloning or subcloning process, and, in particular, for constructing expression vectors by using a site-specific recombinase in vitro or in vivo. In one aspect, a method is used to

directly fuse a linear segment of donor DNA (e.g., gene of interest) with another linear segment of DNA comprising various functional elements such as promoters, selectable markers and a recombination site, resulting in a single circular donor DNA. This circular donor DNA is then
5 recombined into a circular acceptor vector which also contains a recombination site through site-specific recombination catalyzed by a recombinase. The recombination product can be used to transform, transfect or transduce various types of host cells, depending on the specific type of acceptor vector used.

10

The circular donor DNA need not carry an origin of replication for propagation in host cells such as bacterial cells. Instead, the circular donor DNA may be produced from directly ligating two or more linear segments of DNA which may be amplified by polymerase chain reaction
15 (PCR). Such a separation and ligation of different segments of DNA allows flexible distribution of different elements among the linear segments. For example, one linear segment may contain the gene of interest amplified from a cDNA library, while the other linear segment contains functional elements essential for subsequent recombination in
20 vitro (or in vivo) and expression in host cells. By using a site-specific recombinase, such as Cre recombinase, this circular DNA can be recombined into any gene-transferring vector without using restriction enzymes as long as the vector carries a recombination site recognized by the recombinase. Further, direct ligation of linear segments of DNA
25 avoids laborious steps of bacterial cloning and facilitates high throughput screening of large library of genetic materials, such as cDNA libraries derived from diseased tissues or cells. In addition, the circular DNA produced by direct ligation of these segments can be free of other undesirable genetic materials such as "junk DNA" derived from a
30 bacterial plasmid that may affect expression, viability or stability of the recombinant vector.

1. Circular Donor DNA

The present invention provides a double-stranded circular donor DNA for transferring a donor DNA sequence into expression vectors.

5 The circular donor DNA comprises: a donor DNA sequence; a donor recombination site; at least one selectable marker, the circular donor DNA not including an origin of replication. The donor DNA sequence may be any gene of interest which is needed to be transferred into an expression vector.

10

The present invention also provides a library of double-stranded circular donor DNAs which comprises: a donor DNA sequence which varies within a library of donor DNA sequences; a donor recombination site; and at least one selectable marker, the circular donor DNA not
15 including an origin of replication.

The circular donor DNA contains a donor DNA segment (either cDNA or genomic DNA), a promoter (e.g. SV40 early gene enhancer), a selectable marker (e.g. *Neo* gene), and a sequence-specific
20 recombinase target site (e.g. a loxP site). The promoter controls expression of the gene of interest and the selectable marker gene when the circular donor DNA is recombined with an acceptor vector and the resulting recombinant vector is introduced into a host cell. The circular donor DNA may further contain a polyadenylation signal for expression
25 in mammalian cells.

The donor DNA sequence may be any deoxyribonucleotide sequence encoding a functional gene or any synthetically generated DNA sequence. For example the donor DNA segment may be a
30 sequence derived from cDNA of a particular gene or one of the members of a cDNA library. The cDNA library may be produced by

converting mRNAs in a sample into double-stranded complementary DNA (cDNA) by using reverse transcriptase (RT) and the Klenow fragment of nucleic acid polymerase I. Depending on the source of mRNA sample, the cDNA library may contain various populations of genes of interest, such as disease genes located in certain tissue or type of cells. The donor DNA may also be a genomic DNA that contains the coding region interrupted with non-coding sequences (introns/intervening sequences). These introns may contain regulatory elements such as enhancers.

10

The circular donor DNA may further comprises a promoter sequence that controls expression of the donor DNA sequence. The promoter may be any array of DNA sequences that interact specifically with cellular transcription factors to regulate transcription of the downstream gene. The promoter may be derived from any organism, such as bacteria, yeast, insect and mammalian cells and viruses. The selection of a particular promoter depends on what cell type is to be used to express the protein of interest. Examples of the promoter include, but are not limited to, *E. coli lac* and *trp* operons, the *lac* promoter, the bacteriophage λ p^L promoter, bacteriophage T7 and SP6 promoters, β -actin promoter, insulin promoter, human cytomegalovirus (CMV) promoter, HIV-LTR (HIV-long terminal repeat), Rous sarcoma virus RSV-LTR, simian virus SV40 promoter, baculoviral polyhedrin and p10 promoter. The promoter may also be an inducible promoter that regulates the expression of downstream gene in a controlled manner, such as under a specific condition of the cell culture. Examples of inducible promoters include, but are not limited to, the bacterial dual promoter (activator/repressor expression system) which regulates gene expression in mammalian cells under the control of tetracyclines (Gossen, M. and Bujard, H. 1992, Proc.Natl. Acad. Sci. USA, 89, 5547-5551) and promoters that regulate gene expression under the control of

30

factors such as heat shocks, steroid hormones, heavy metals, phorbol ester, the adenovirus E1A element, interferon, or serum.

5 The recombination site may be any segment or arrays of DNA sequence recognized by site-specific recombinase which catalyzes site-specific fusion between the circular donor DNA and the acceptor vector. The site-specific recombinase may be any enzymes that recognize short DNA sequences that become the crossover regions during the recombination event, including but not limited to recombinases,
10 transposases and integrases.

Site-specific recombinases may derived from prokaryotic and eukaryotic sources. Examples of site-specific recombination include 1) chromosomal rearrangements which occur in *Salmonella typhimurium*
15 during phase variation, inversion of the FLP sequence during the replication of the yeast 2 μ circle and in the rearrangement of immunoglobulin and T cell receptor genes in vertebrates, 2) integration of bacteriophages into the chromosome of prokaryotic host cells to form a lysogen and 3) transposition of mobile genetic elements (e.g.,
20 transposons) in both prokaryotes and eukaryotes.

In one embodiment, the recombination site is a loxP site that is recognized by the Cre recombinase of bacteriophage ϕ I. The Cre recombinase catalyzes recombination of DNA between two loxP sites.
25 The loxP site consists of a double-stranded 34 bp sequence:

5'-ATAACTTCGTATAATGTATGCTATACGAAGTTAT-3'
3'-TATTGAAGCATATTACATACGATATGCTTCAATA-5'

30 (SEQ ID NO:1)

The loxP site comprises two 13 bp inverted repeat sequences separated by an 8 bp spacer region. Hoess et al. (1982) Proc. Natl. Acad. Sci. USA 79:3398. The internal spacer sequence of the loxP site is asymmetrical and thus, two loxP sites can exhibit directionality relative to one another. Hoess et al. (1984) Proc. Natl. Acad. Sci. USA 81:1026. When two loxP sites on the same DNA molecule are in a directly repeated orientation, Cre excises the DNA between these two sites leaving a single loxP site on the DNA molecule. Abremski et al. (1983) Cell 32:1301. If two loxP sites are in opposite orientation on a single DNA molecule, Cre inverts the DNA sequence between these two sites rather than removing the sequence.

The Cre recombinase also recognizes a number of variant or mutant lox sites relative to the loxP sequence. Examples of these Cre recombination sites include, but are not limited to, the loxB, loxL and loxR sites which are found in the E. coli chromosome. Hoess et al. (1982), supra. Other variant lox sites include:

loxP511 site: 5'-ATAACTTCGTATAGTATACATTATACGAAGTTAT-3'
(SEQ ID NO:2);
Hoess et al. (1986) Nucleic Acid Res. 14:2287-2300,

loxC2 site: 5'-ACAAC TTCGTATAATGTATGCTATACGAAGTTAT-3'
(SEQ ID NO:3)
U.S. Pat. No. 4,959,317.

Cre catalyzes the cleavage of the lox site within the spacer region and creates a six base-pair staggered cut. Hoess and Abremski (1985) J. Mol. Biol. 181:351. The two 13 bp inverted repeat domains of the lox site represent binding sites for the Cre protein. If two lox sites differ in their spacer regions in such a manner that the overhanging ends of the

cleaved DNA cannot reanneal with one another, Cre may not efficiently catalyze a recombination event using the two different lox sites. For example, it has been reported that Cre could not efficiently recombine a loxP site and a loxP511 site; these two lox sites differ in the spacer
5 region. Two lox sites which differ due to variations in the binding sites (i.e., the 13 bp inverted repeats) may be recombined by Cre provided that Cre can bind to each of the variant binding sites; the efficiency of the reaction between two different lox sites (varying in the binding sites) may be less efficient than between two lox sites having the same
10 sequence (the efficiency will depend on the degree and the location of the variations in the binding sites). For example, the loxC2 site can be efficiently recombined with the loxP site; these two lox sites differ by a single nucleotide in the left binding site.

15 The Cre protein has been purified to homogeneity. Abremski et al. (1984) J. Mol. Biol. 259:1509. And the *cre* gene has been cloned and expressed in a variety of host cells. Abremski et al. (1983), *supra*. Purified Cre protein is available from a number of suppliers (e.g., Novagen and New England Nuclear/Du Pont).

20 The recombination site of the circular DNA may also be selected from a variety of other recombination sites recognized by recombinases other than Cre. Examples of the non-Cre recombinases include, but are not limited to, site-specific recombinases include: the Int recombinase
25 of bacteriophage λ , the FLP recombinase of the 2 π i plasmid of *Saccharomyces cerevisiae*, the resolvase family, transposase of *Bacillus thuringiensis*.

30 The Int recombinase of bacteriophage λ belongs to the integrase family and mediates the integration of the λ genome into the *E. coli* chromosome. The Int recombinase of bacteriophage λ promotes

irreversible recombination between its substrate att sites as part of the the formation or induction of a lysogenic state. Landy, A., (1989) Ann. Rev. Biochem. 58:913. Reversibility of the recombination reactions results from two independent pathways for integrative and excisive recombination. Each pathway uses a unique but overlapping set of the 15 protein binding sites that comprise att site DNAs. Cooperative and competitive interactions involving four proteins (Int, Xis, IHF and FIS) determine the direction of recombination. Integrative recombination involves the Int and IHF proteins and sites attP (240 bp) and attB (25 bp). Recombination results in the formation of two new sites: attL and attR. Excisive recombination requires Int, IHF, and Xis, and sites attL and attR to generate attP and attB. Under certain conditions, FIS stimulates excisive recombination. In addition to these normal reactions, it should be appreciated that attP and attB, when placed on the same molecule, can promote excisive recombination to generate two excision products, one with attL and one with attR. Similarly, intermolecular recombination between molecules containing attL and attR, in the presence of Int, IHF and Xis, can result in integrative recombination and the generation attP and attB. Derivatives of the att site with changes within the 15 bp core may also be suitable for efficient recombination. By incorporating a native or modified att site in both the circular donor DNA and the acceptor vector, intermolecular recombination between the donor and acceptor DNA molecules may be achieved by using the appropriate recombination protein such as Int, IHF and FIS, with or without Xis. Integrase can be obtained as described by Nash, H. A., (1983) Methods of Enzymology 100:210-216. IHF can be obtained as described by Filutowicz, M., et al., (1994) Gene 147:149-150.

The other members of the integrase family of site-specific recombinases may also be used to provide alternative recombination proteins and recombination sites for the present invention. Examples of

such Int recombinases include, but not limited to, site-specific recombinase encoded by bacteriophage λ , phi 80, P22, P2, 186, P4.

This group of recombinases exhibits a large diversity of sequences, but all of the recombinases can be aligned in their C-terminal halves. A 40-residue region near the C terminus is particularly well conserved in all the proteins and is homologous to a region near the C terminus of the yeast 2 μ plasmid Fip protein. Three positions are perfectly conserved within this family: histidine, arginine and tyrosine are found at respective alignment positions 396, 399 and 433 within the well-conserved C-terminal region. These residues contribute to the active site of this family of recombinases, and suggest that tyrosine-433 forms a transient covalent linkage to DNA during strand cleavage and rejoining. Argos, P. et al., (1986) EMBO J. 5:433-40.

The FLP recombinase of the 2 μ plasmid of *Saccharomyces cerevisiae* recognizes the *frt* site which, like the *loxP* site, comprises two 13 bp inverted repeats separated by an 8 bp spacer:

5'-GAAGTTCCTATTCTCTAGAAAGT ATAGGAACTTC-3'

(SEQ ID NO:4)

Cox (1983) Proc. Natl. Acad. Sci. USA 80:4223.

The FLP gene has been cloned and expressed in *E. coli* and in mammalian cells and has been purified. Meyer-Lean et al. (1987) Nucleic Acids Res. 15:6469; Babineau et al (1985) J. Biol. Chem. 260:12313; Gronostajski and Sadowski (1985) J. Biol. Chem. 260:12328.

The resolvase family members, such as the Tn3 resolvase, the Hin recombinase, and the Cin recombinase, may also be used for recombination between the circular donor DNA and the circular acceptor

DNA. Although members of this highly related family of recombinases are typically constrained to intramolecular reactions (e.g., inversions and excisions) and can require host-encoded factors. Mutants have been isolated that relieve some of the requirements for host factors as well as
5 some of the constraints of intramolecular recombination. Maeser and Kahnmann (1991) Mol. Gen. Genet. 230:170-176.

Transposase of *Bacillus thuringiensis* may also be used as recombination proteins and recombination sites. *Bacillus thuringiensis* is
10 an entomopathogenic bacterium whose toxicity is due to the presence in the sporangia of Δ -endotoxin crystals active against agricultural pests and vectors of human and animal diseases. Most of the genes coding for these toxin proteins are plasmid-borne and are generally structurally associated with insertion sequences (IS231, IS232, IS240, ISBT1 and
15 ISBT2) and transposons (Tn4430 and Tn5401). Several of these mobile elements have been shown to be active and participate in the crystal gene mobility, thereby contributing to the variation of bacterial toxicity. Structural analysis of the iso-IS231 elements indicates that they are related to IS1151 from *Clostridium perfringens* and distantly related to
20 IS4 and IS186 from *E. coli*. Like the other IS4 family members, they contain a conserved transposase-integrase motif found in other IS families and retroviruses. Functional data gathered from IS231A in *E. coli* indicate a non-replicative mode of transposition, with a preference for specific targets. Similar results were also obtained in *Bacillus subtilis*
25 and *B. thuringiensis*. Mahillon, J. et al., (1994) Genetica 93:13-26; (1992) Campbell, J. Bacteriol. 74:95-7499.

Other recombination systems may also be used as recombination proteins and recombination sites, including the xerC and xerD
30 recombinases of *E. coli* which together form a recombinase that recognizes the 28 bp dif site (Leslie and Sherratt (1995) EMBO J.

14:1561); the Int protein from the conjugative transposon Tn916 (Lu and Churchward (1994) EMBO J. 13:1541); TpnI and the β -lactamase transposons (Levesque (1990) J. Bacteriol. 172:3745); and the immunoglobulin recombinases (Malynn et al. (1988) Cell 54:453).

5

Other than the wild-type recombination sites, modified recombination sites may also be used in the present invention. Wild-type recombination sites may contain sequences that reduce the efficiency or specificity of recombination reactions. For example, multiple stop
10 codons in attB, attR, attP, attL and loxP recombination sites occur in multiple reading frames on both strands, thereby reducing recombination efficiencies. For example att sites, such as att1, att2, and att3 sites, may be modified to have one or multiple mutations to enhance specificity or efficiency of the recombination reaction and to
15 decrease reverse reaction by removing P1 and H1 from attB.

The circular donor DNA also contains one or more selectable markers to facilitate subsequent identification and selection of clones of the recombination product under suitable conditions. The selectable
20 marker may encode any functional element, such as protein, peptide, RNA, binding site for RNA and proteins, or products that provide resistance to organic or inorganic agents. Examples of selectable markers include, but are not limited to, reporter genes such as β -galactosidase (GAL), fluorescent proteins (e.g., GFP, GFP-UV, EFP, BFP, EBFP, ECFP, EYFP), secreted form of human placental alkaline
25 phosphatase (SEAP), β -glucuronidase (GUS)); resistance genes that encode products which provide resistance against otherwise toxic agents such as antibiotics (e.g. neomycin (G418) or hygromycin resistant gene, puromycin resistant gene), yeast selectable markers leu2-
30 d and URA3, apoptosis resistant genes (e.g. the baculoviral p35 gene) that encode proteins that binds to products which are detrimental to cell

survival and promote apoptosis; antisense oligonucleotides, and any other DNA that encodes product that directly or indirectly confer sensitivity of cells to particular agents.

5 The circular donor DNA may optionally include an affinity tag for selection and isolation of protein product encoded by the donor DNA segment. Examples of such an affinity tag include, but are not limited to, a polyhistidine tract, polyarginine, glutathione-S-transferase (GST), maltose binding protein (MBP), a portion of staphylococcal protein A
10 (SPA), and various immunoaffinity tags (e.g. protein A) and epitope tags such as those recognized by the EE (Glu-Glu) antipeptide antibodies. The affinity tag may also be a signal peptide either native or heterologous to baculovirus, such as honey bee mellitin signal peptide. The affinity tag may be positioned at either the amino- or carboxy-terminus of the
15 donor DNA.

2. Circular Acceptor Vector

20 The present invention also provides a circular acceptor vector for generating recombinant expression vector. The vector comprises an origin of replication; and an acceptor recombination site capable of recombining with a donor DNA. Optionally, the acceptor vector may not include a promoter for regulating expression of the donor DNA.

25 The circular acceptor vector may be any vector that can transform, transfect or transduce a host cell. The acceptor vector comprises a recombination site which is recognized by a site-specific recombinase and recombined with a donor DNA carrying another recombination site. The acceptor vector may be plasmids, phages or
30 viral vectors as long as it is able to replicate in vitro, or in a host cell, or to convey the donor DNA to a desired location within a host cell.

Examples of host cells include, but are not limited to, bacterial (e.g. *E. coli*, *Bacillus subtilis*, etc.), yeast, animal, plant, and insect cells.

5 For plasmid-based expression vector, the recombination site may be introduced into the vector by a double-stranded oligonucleotide containing the desired site-specific recombination site (e.g., a *lox* site). The double-stranded oligonucleotide may be formed by annealing two synthetic single-stranded oligonucleotides to form two ends which are compatible with ends of a linearized plasmid vector. The matching ends
10 may be generated by restriction enzyme digestion or by using cloning kits such as the TA cloning kits available from Invitrogen, Inc. (San Diego, CA).

The circular acceptor vector may be any prokaryotic plasmid that
15 contains a recombination site (e.g. *loxP* site), a basic backbone of plasmid cloning vector such as pBR322, including one or more antibiotic resistant genes (e.g. *Amp^r*, *Tet^r*) and an origin of replication that function in specific host cells. After recombination between the circular donor DNA and the acceptor vector to form a fused plasmid, this plasmid
20 vector can be used to transform bacterial cells. In the transformed cell, a prokaryotic promoter, either carried by the donor DNA or the acceptor vector, causes expression of the donor DNA under suitable conditions.

Optionally, the acceptor vector may comprise a prokaryotic
25 termination sequence. Examples of the prokaryotic termination sequence include, but are not limited to, the T7 termination sequence. A variety of termination sequences are known to the art and may be employed in the nucleic acid constructs of the present invention including, the T_{INT} , T_{L1} , T_{L2} , T_{L3} , TR_1 , TR_2 , T_{6S} termination signals derived
30 from the bacteriophage ϕ . Hendrix et al. Eds., Cold Spring Harbor Press, Cold Spring Harbor, N.Y. (1983) and termination signals derived

from bacterial genes such as the trp gene of E. coli.

5 The circular acceptor vector may also be a mammalian expression vector that contains a recombination site (e.g. loxP site), one or more eukaryotic marker gene, the appropriate eukaryotic transcriptional and translational termination signals and a sequence that signals polyadenylation of the transcript messenger RNA (mRNA), and an origin of replication that functions in a mammalian host cell. Examples of the eukaryotic polyadenylation sequence include, but are not limited to, the Herpes simplex virus thymidine kinase polyadenylation sequence, the bovine growth hormone polyadenylation sequence, and the simian virus 40 polyadenylation sequence.

15 If the circular acceptor carries a promoter for regulating expression of the donor DNA sequence, the recombination site may be placed downstream the promoter and transcription initiation site in acceptor vector. This modification of the vector may be easily accomplished using synthetic oligonucleotides comprising the desired recombination site (loxP site). In designing the oligonucleotide comprising the recombination site, it may be desirable to avoid introducing an ATG or start codon that might initiate translation inappropriately, or in-frame stop codons.

25 For expression vectors intended to generate a fusion protein between a protein domain located at the amino-terminus of the fusion protein and the protein encoded by the donor DNA, care may be taken to place the recombination site in the correct reading frame such that 1) an open reading frame is maintained through the recombination site on pHOST and 2) the reading frame in the recombination site on the acceptor vector is in frame with the reading frame found on the recombination site contained within the circular donor DNA.

Optionally, the eukaryotic expression vector may also carry an origin of replication and selectable marker genes that function in bacterial cells, forming a shuttle vector. After recombination between the circular donor DNA and the acceptor vector to form a fused expression vector, this vector can be used to transfect animal cells. In the transfected cell, a eukaryotic promoter, either carried by the donor DNA or the acceptor vector, causes expression of the donor DNA under suitable conditions.

The circular acceptor may also be a yeast expression vector such as a *S. cerevisiae* expression vector that includes a recombination site. Various types of *S. cerevisiae* expression vector include episomal or plasmid vector, integrating vectors, and yeast chromosomes (YACs).

A YAC-based expression vector may be used to carry large segment of donor DNA, which is then maintained as a separate chromosome in the host yeast cell.

The circular acceptor vector may also be a baculovirus DNA (genome) that is modified to contain a recombination site (e.g. a loxP site). For example, the baculoviral genome of *Autographa californica* multiple nuclear polyhedrosis virus (AcMNPV) may be modified to include a site-specific recombination site, such as a loxP site. *A. californica* (the alfalfa loop) and over 30 other insect species can be infected by AcMNPV. This virus also grows well on many insect cell lines, such as Sf cell lines derived from the fall armyworm, *Spodoptera frugiperda*. In these cells, the the promoter of the viral protein, polyhedrin, is exceptionally strong when the virus infects the cell. Such polyhedrin promoter can promote high level expression of downstream foreign gene that replaces the polyhedrin gene.

The recombination site (e.g. a loxP site) may be introduced into a baculoviral vector through homologous recombination in vivo. For example, a baculoviral transfer vector may be used to shuttle the loxP site into the wild type baculoviral DNA (e.g. AcMNPV) to form the final circular baculoviral vector. An example of such transfer vector is the p36C that contains baculoviral polyhedrin flanking sequences, up- and down-stream AcMNPV DNAs for homologous recombination. Page (1989) Nucleic Acids Res. 17: 454. The loxP site may be introduced into the transfer vector by regular oligonucleotide-mediated mutagenesis. A portion of the polyhedrin flanking sequence is replaced with the loxP site using an oligonucleotide containing the loxP site. The resulting loxP-containing transfer vector and the wild type baculoviral DNA are co-transfected into insect cells such as Sf9 cells. Because the transfer vector contains the polyhedrin flanking sequences, a double crossover homologous recombination occurs in the cells, causing the replacement of the polyhedrin gene in AcMNPV DNA with the loxP site and therefore resulting in the integration of the loxP site into the AcMNPV genome. After a desired period of time (e.g. 72 hr) the supernatant of the insect cell culture is harvested and the progeny virus is screened in a standard agarose overlay assay. Brown and Faulkner (1977) J. Gen. Virol. 36: 361-364. Polyhedrin-negative plaques are purified to homogeneity by successive rounds of agarose overlay assay. The presence of the loxP site may be confirmed by Southern analysis of DNA from the infected cells.

Optionally, the baculoviral acceptor vector according to the present invention may not contain the polyhedrin promoter. Instead, the polyhedrin promoter or a baculoviral p10 promoter is positioned upstream of the donor DNA sequence of the circular donor DNA. After site-specific recombination between the circular donor DNA and a baculoviral acceptor vector, the resulting recombinant baculoviral vector

can be used to infect insect cells, such as Sf9 cells.

Also optionally, the circular acceptor vector according to the present invention may include a negatively selectable marker gene,
5 such as the selectable marker based on herpes simplex virus tk gene or any kind of gene encoding a protein that signals apoptosis and causes programmed cell death, such as the CAR1 gene. Expression of these negatively selectable marker genes results in cell death, thereby eliminating those cells containing the circular acceptor vector that does
10 not recombine with the circular donor DNA.

The present invention also provides kits for generating recombinant vectors. In one embodiment, the kit comprises: a double-stranded circular donor DNA comprising a donor DNA sequence, a
15 donor recombination site, and at least one selectable marker, the circular donor DNA not including an origin of replication; and a circular acceptor vector comprising an origin of replication and an acceptor recombination site capable of recombining with the circular donor DNA.

20 In another embodiment the kit comprises: a library of double-stranded circular donor DNA comprising a donor DNA sequence which varies within a library of donor DNA sequences, a donor recombination site, and at least one selectable marker, the circular donor DNA not including an origin of replication; and a circular acceptor vector
25 comprising an origin of replication and an acceptor recombination site capable of recombining with the circular donor DNA.

In yet another embodiment, the kit comprises: one or more linear donor DNA comprising a donor DNA sequence; a linear driver DNA
30 comprising a promoter sequence, a recombination site, and at least one selectable marker, ligation of the linear donor DNA and the linear driver

DNA resulting in a circular donor DNA; and a circular acceptor vector comprising an origin of replication and an acceptor recombination site capable of recombining with the circular donor DNA.

5

3. Method of Generating Recombinant Expression Vectors through Site-Specific Recombination

The present invention provides a method for generating
10 recombinant expression vector *in vitro* through site-specific recombination between a circular donor DNA and circular acceptor DNA, each containing recombination site recognized by the recombinase. The method comprises: contacting a circular double-stranded donor DNA and a circular acceptor vector in the presence of a
15 recombinase under conditions suitable for the circular double-stranded donor DNA and circular acceptor vector to recombine to form a single fused circular vector.

In this method, the circular double-stranded donor DNA
20 comprises a donor DNA sequence, a donor recombination site, and at least one selectable marker, but not including an origin of replication. The circular acceptor vector comprises an origin of replication and an acceptor recombination site capable of recombining with the circular donor DNA. The promoter for regulating expression of the donor DNA
25 may be contained in either the donor DNA or acceptor vector.

According to this method, the circular donor DNA containing a site-specific recombination site can be recombined with a circular acceptor vector in the presence of a site-specific recombinase, such as
30 Cre recombinase. *In vitro* Cre recombinase can catalyze fusion of the donor DNA into the acceptor vector which contains another copy of the

recombination site (e.g. loxP site) found on the circular donor DNA. Optionally, the circular donor DNA may contain a promoter positioned upstream of the donor DNA sequence to regulate expression of the donor DNA once the recombinant vector is introduced into a host cell.

5

Alternatively, the promoter for regulating expression of the donor DNA sequence may be positioned upstream of the recombination site of the acceptor vector. Following the site-specific recombination between the recombination sites located on the donor DNA and acceptor vector, the two circular DNAs are stably fused in a manner that places the expression of the donor DNA sequence under the control of the promoter contained within the acceptor vector. The recombination occurs in a manner that retains the proper translational reading frame of the donor DNA.

15

Following the in vitro recombination, a portion of the reaction mixture may be used to transform, transfect or transduce a suitable host cell to permit the recovery and propagation of the recombinant expression vectors. The correctly fused recombinant vector may be selected for its ability to transform, transfect or transduce a host cell and express the selectable marker that is contained in the circular donor DNA and recombined into the acceptor vector. The selectable phenotype conferred by the selectable marker gene on the recombinant vector may be change of color of the host cells upon proper chemical treatment, secretion of protein in the culture that is detectable by colorimeters or fluoreimeters, survival of cells under selection pressure or enhanced propagation of a virus such as baculovirus. The circular donor DNA cannot replicate in cells because it does not contain an origin of replication, and therefore, unless the circular donor DNA has integrated into the acceptor vector that contains an origin of expression, the circular donor DNA should not replicate in the host cell. The

20
25
30

recombinant expression vector may be isolated from host cells which display the desired phenotype and verified by using various methods such as restriction enzyme digestion, sequencing and Southern blotting.

5 The present invention also provides a method for generating recombinant expression vectors from linear DNA segments *in vitro*. The method comprises: ligating one or more double-stranded linear donor DNA which includes a donor DNA sequence with a double-stranded linear driver DNA which includes a promoter sequence and a donor
10 recombination site to form a single circular donor DNA, the singular circular donor DNA not including an origin of replication, where the donor DNA sequence is under the transcriptional control of the promoter; and contacting the circular donor DNA and a circular acceptor vector in the presence of a recombinase to form a single fused
15 circular vector. In this method, the circular acceptor vector comprises an origin of replication and an acceptor recombination site capable of recombining with the circular donor DNA.

 According to this method, the linear donor DNA and linear driver
20 DNA may contain matching restriction sites or other type of annealing sites so as to be ligated to form a circularized DNA. After the ligation, the circularized donor DNA contains a site-specific recombination site carried by the linear driver DNA. Such a circular donor DNA is recombined with a circular acceptor vector in the presence of a site-
25 specific recombinase, such as Cre recombinase. The Cre recombinase can catalyze fusion of the donor DNA into the acceptor vector which contains another copy of the recombination site (e.g. loxP site) found on the circular donor DNA.

30 Figure 1 illustrates a general scheme of this method according to the present invention. As illustrated in Figure 1, a library of linear donor

DNA (underlined) is directly ligated to a linear driver DNA (underlined) to generate a circular donor DNA. This allows for efficient cloning of donor DNA (e.g. cDNA library) into an expression vector in an automated and high throughput manner.

5

The linear donor and driver DNA can be generated by PCR amplification of a template DNA, either linear or contained in plasmids of cDNA clones. As illustrated in Figure 1, a library of cDNA clones from a desired source may be rearrayed into 96-well plates by using a re-
10 arraying robot. The library of cDNA clones may be cosmid and phage libraries that contain inserts primarily from single human chromosomes isolated by flow-sorting. These cDNA clones may be identified by their short sequence tags or expressed sequence tags (ESTs). These cDNA clones may be arrayed in multi-well microtiter plates such as 96- or 384-
15 well plates and handled by robots. For each re-arrayed plate, data tracking system may be used to identify each clone, sequence accession number, passage number, etc.

The library of linear donor DNA may also be generated and
20 amplified from total RNA or mRNA samples by using an RT-PCR method. The library of linear donor DNA can then be ligated with the linear driver DNA to form the circular donor DNA which is recombined with the circular acceptor. This allows direct transferring of the cDNA library (donor DNA) to the expression vector (acceptor vector) without
25 going through a cloning step.

Alternatively, the library of linear donor DNA may be generated by random or site-directed mutagenesis of one or more target gene sequence. For example, "poisoned" PCR or DNA shuffling techniques
30 may be used to generate a diverse library of donor DNA which can be incorporated into the circular donor DNA by using direct ligation

according to the present invention. Clones of these mutagenized library can be screened for improved or unique biological functions in an appropriate expression system.

5 Bacterial cultures of the cDNA clones may be grown in the multi-well microtiter plates and directly used for PCR. Suitable PCR primers containing an annealing site may be used to amplify each cDNA clone in the plate. For example, PCR primers may be designed to hybridize to the sequence of the cDNA clones and contain sequences for Ligation
10 Independent Cloning (LIC). High-fidelity thermostable polymerases are preferably used to reduce copying errors and numbers of amplification cycles. The PCR product, the linear cDNA library, may optionally be purified by passing through mini columns.

15 Similarly, the linear driver DNA may also be amplified by using primers containing a matching annealing site as the primers for amplifying donor DNA. The template of the driver DNA may also be a plasmid which contains functional elements, such as a site-specific recombination site, a promoter, a selectable marker gene, a tag, and a
20 transcription termination signal.

 Still referring to Figure 1, the linear driver DNA is annealed and ligated with the linear donor DNA under suitable conditions in multiple-well plates (e.g. in the presence of T4 ligase). For ligation independent
25 cloning (LIC), the linear driver DNA and the linear donor DNA generated by LIC PCR amplification are digested with T4 polymerase, annealed, and ligated together to form a library of circular donor DNAs
 (underlined).

30 The circular donor DNA may be mixed with a circular acceptor vector (underlined) containing a suitable recombination site (e.g. loxP) in

the presence of a site-specific recombinase (e.g. Cre recombinase). For example, a circular baculoviral vector containing a loxP site may be mixed with an array of circular cDNA (circular donor DNA) in the presence of purified Cre recombinase or cell extract containing Cre.

- 5 The amount of recombinase which is added to drive the recombination reaction between the donor and vector DNAs can be determined by using a titration assay is used to determine the appropriate amount of the purified recombinase enzyme or the appropriate amount of a Cre extract. The recombination reaction buffer compatible with purified Cre
- 10 recombinase may contain 50 mM Tris-HCl (pH 7.5), 10 mM MgCl₂, 30 mM NaCl and 1 mg/ml BSA. The concentration of the circular donor DNA and the acceptor vector may vary between 10 ng to 10 µg of each vector per 20 µl reaction volume. The recombination reaction may be incubated at 37°C for a necessary period of time and terminated by
- 15 heating at 70°C for 15 min.

Site-specific recombination between the donor cDNA array and the acceptor baculoviral vector in the presence of Cre recombinase results in a recombinant baculoviral vector (the vector DNA in Figure 1).

- 20 The recombinant baculoviral vector may be used to transfect Sf9 cells (the host cell in Figure 1) arrayed in multi-well plates. The plates may be incubated at 27°C for several days. The progeny baculovirus may be screened in an overlay assay of the supernatant. For example, the presence of the integrated donor DNA in the viral progeny may be
- 25 determined by detecting level of expression of the selectable marker (e.g. β-GAL, GUS). Alternatively, a baculovirus apoptosis resistant gene (baculovirus p35 gene) may be used as a selectable marker to confer increased viral yield in certain insect cell lines. Clem et al. (1991) Science 254: 1388-1390. Recombinant baculovirus bearing this gene
- 30 has been shown to be amplified up to a million folds in appropriate host cells. Lerch et al (1993) Nucleic Acid Res. 21: 1753-1760.

The protein product expressed by the recombinant expression vector may be selectively identified and purified by using affinity tagging. The tag element may be a short DNA sequence inserted between the promoter and the transcription termination sequence in a proper reading frame on the recombinant expression vector. Such a tag element may encodes a short amino acid sequence that specifically binds to a compound or a macromolecule. After transcription and translation of the recombinant expression vector in the host cells the short amino acid sequence which is part of the donor DNA construct, acts as an identification tag (affinity tag). In another word, the tag carried by the circular donor DNA that is recombined with the acceptor DNA is expressed as a fusion protein encoded by the donor DNA. The presence of the fusion protein can be identified through the affinity binding of the tag to its binding partner.

For example, a polyhistidine tag (e.g. His₆) may be used for isolating fusion proteins. The fusion protein carrying a His tag can be isolated by passing all proteins of a cell extract (e.g. Sf9 cells) through a column of Ni-triacetic acid agarose beads. The unbound proteins are eluted from the column, and the bound fusion protein is selectively removed by either adding a competitor compound (e.g. imidazole) that dislodges the His tag of the fusion protein from the Ni ions or by lowering the pH of the elution buffer. If necessary, the tag may be removed from the fusion protein with a protease that cleaves only at the engineered site.

Alternatively, the affinity tag may be a protein tag (e.g., GST, maltose binding protein) or an epitope tag that are short amino acid sequences binding to glutathione, maltose and specific antibodies. For example, an antibody against an EE epitope tag may be used for

identifying and purifying fusion protein encoded by the donor DNA. The EE (Glu-Glu) antipeptide antibody was raised against a peptide that contains the major tyrosine phosphorylation site of polyoma middle T antigen. Talmage et al. (1989) Cell 59: 55-56. The antibody has moderate affinity for the EE tag ($K_d \sim 2 \times 10^{-7}$) which allows rapid elution of tagged fusion proteins by free peptide under non-denaturing conditions while retaining efficient binding of the fusion protein in crude lysates. Further, since the EE tag is a strong tyrosine kinase phosphorylation substrate for protein kinases (e.g. Src) it may serve as a detection label for high throughput assays for protein interactions.

The methods of the present invention allow rapid and efficient generation of expression vectors containing the gene of interest without bacterial cloning. Various libraries of cDNA or genomic DNA that are difficult to be cloned in bacteria can be directly amplified and introduced into any expression vector. The methods can be used in a wide variety of high through arrays for functional genomics, protein genomics and other human genome projects. For example, the methods may be used for systematic functional analysis gene profiling, gene tagging, gene overexpression, or systematic transcript analysis. The information generated can shed light on functionally important pathways in diseased cells in many important areas such as oncology and inflammation. Further, the methods may be used for high-throughput genetic screens for target discovery and validation, as well as drug discovery based on the targets discovered in the screens.

EXAMPLE

Figure 2 illustrates an example of how to construct a baculoviral expression vector for the GUS gene according to the present invention. As illustrated in Figure 1, a recombinant baculoviral expression vector is

generated *in vitro* without a cloning step. The following is a detailed elucidation of the steps of using the method.

1. Construction of the driver plasmid

5

A linear double-stranded driver DNA (underlined) is generated from a driver plasmid containing a baculoviral p10 promoter, a *Neo* marker gene, a loxP site and appropriate restrictions sites such as Sfi I and Not I sites. The Driver plasmid is constructed by using standard plasmid construction techniques as taught in Sambrook, et al.

10

Molecular Cloning: A laboratory Manual; *DNA Cloning: A Practical Approach*, Vol I & II (D. Glover ed.); *Oligonucleotide Synthesis* (N. Giat, ed.). The driver plasmid is based on pBR322, a standard bacterial cloning vector plasmid, and the following elements are cloned into pBR322 contiguously, in the order shown below:

15

- a) an SfiI restriction site (GGCCNNNNNGGCC)
- b) a loxP site
(ATAACTTCGTATAATGTATGCTATACGAAGTTAT)
[SEQ ID NO: 1]
- c) a *Neo* gene with promoter and SV40 poly A addition site
- d) a baculovirus p10 promoter
- e) an EE-tag sequence (MEEEEYMPME) [SEQ ID NO: 5]
- f) a NotI restriction site (GCGGCCGC)

20

25

Sequence of the baculoviral p10 promoter is described in Weyer U, Possee RD. (1989) J. Gen. Virol. 70:203-8 and is supplied by pAcAB4 from BD Pharmingen, San Diego, CA. The sequence for the *Neo* gene with a promoter and SV40 poly A addition site is available from the plasmid pLE-neo supplied by Novagen Inc.

30

2. Amplification and processing of the donor gene DNA

A linear double-stranded donor DNA (underlined) is amplified by PCR from a donor plasmid or a cDNA clone carrying the GUS gene.

5 Oligonucleotide PCR primers that are homologous to the 5' and 3' ends of the GUS gene (source: pBacPAK8-GUS available from Clontech, Palo Alto, CA) are designed and synthesized. These primers also contain restriction sites (NotI for the 5' primer and SfiI for the 3' primer) in the 5' ends of the primers. The GUS gene is then amplified by PCR
10 from the plasmid pBacPAK8-GUS containing the GUS gene using these primers.

The PCR products from GUS gene amplification are digested with NotI and SfiI restriction endonucleases to generate the appropriate
15 sticky ends on the products, resulting in the linear double-stranded donor DNA.

3. Construction of the circular donor DNA

20 The driver plasmid described in Section 1 is digested with NotI and SfiI restriction endonucleases. The fragment which contains the elements listed above such as the loxP site and p10 promoter is isolated by gel electrophoresis purification, resulting in the linear double-stranded driver DNA.

25 The linear driver DNA is annealed to an excess of the linear donor DNA and ligated by using T4 DNA ligase to produce the circular donor DNA (underlined).

4. Construction of the acceptor baculovirus genomic DNA:

30 A baculovirus transfer vector plasmid containing a loxP site is constructed according to the protocol described by Peakman et al.

(1992) Nucleic Acid Res. 20:495-500. Briefly, a double-stranded oligonucleotide consisting of a loxP sequence [SEQ ID NO:1] with a blunt 5' end and a CTAG 5' overhang at the 3' end is synthesized. A baculovirus transfer vector pVL1392 (available from BD Pharmingen, San Diego, CA) is digested with EcoR V and BamH I restriction enzymes to remove the polyhedrin promoter in the vector. The double-stranded oligonucleotide with the loxP site is ligated with the EcoR V/BamH I digested plasmid pVL1392. The resulting ligated plasmid is transformed into bacteria and screened for colonies with the loxP insert.

10 This generates a baculoviral transfer vector containing a loxP site.

The baculoviral transfer vector containing a loxP site is co-transfected with a linearized baculovirus DNA (BacPAK6 DNA from Clontech, Palo Alto, CA) by using standard baculovirus construction techniques (Methods in Molecular Biology Vol. 39: Baculovirus Expression Protocols, Christopher D. Ricardson ed., Humana Press, Totowa, NJ 1995).

Recombinant baculovirus that contains the loxP site is isolated by plaque purification and baculoviral DNA is prepared from the virus to produce the circular acceptor DNA.

20

5. Recombination of the circular donor DNA and the circular acceptor DNA and selection of the recombinant baculoviral DNA

The circular acceptor DNA and the circular donor DNA generated as described above is recombined in the presence of GST-Cre recombinase (available from Invitrogen, San Diego, CA) by using the methods described by Liu et al. (1998) Current Biology 8:1300-1309. The mixture is then heated at 65°C to inactivate the Cre recombinase.

25 The circular acceptor DNA that has recombined with the circular donor DNA is selected and purified from the recombination reaction

30

mixture by using a method of "physical selection" described in Shepard & Rae (1997) Nucleic Acid Research 25:3183. Briefly, a biotinylated synthetic oligonucleotide probe that is substantially homologous to any region present in the circular donor DNA is used to hybridize with the recombined baculoviral DNA. The hybridized complex is then separated by using avidin-coated magnetic beads via biotin-avidin high affinity binding. The circular donor DNA that is not recombined with the acceptor DNA may also be extracted from the mixture. Since the circular donor DNA is incapable of replicating in a host insect cell and thus does not interfere with further functional analysis of the recombined baculoviral vector. The biotinylated oligonucleotide probe is preferably homologous to a GC-rich region of the circular donor DNA.

6. Propagation and selection of the recombinant baculovirus

The recombinant baculovirus DNA selected in the above-described process is transfected into Sf9 insect cells using standard baculovirus construction techniques (Methods in Molecular Biology Vol. 39: Baculovirus Expression Protocols, Christopher D. Ricardson ed., Humana Press, Totowa, NJ 1995).

The recombinant GUS baculovirus is passaged in the presence of G418 as described by Lerch and Friessen (1993) Nucleic Acid Res. 21:1753-1760. This is a positive selection step for propagation of the recombinant GUS baculovirus containing the *Neo* gene as a selectable marker.

7. Assessment of recombinant protein production

The GUS activity is assayed by using the following protocol. The recombinant GUS baculovirus produced as described above is used to

infect Sf9 insect cells. After 2 days the Sf9 cells are lysed in a solution of: 20mM Tris pH8, 150 mM NaCl, 0.5% NP40 (10ml of solution per gram of cells). The resulting lysate is serially diluted into PBS plus 150 ug/ml x-Gluc. Blue color in diluted samples indicates GUS activity. The
5 terminal dilution of the recombinant GUS baculovirus produced by using the method of the present invention is compared with that of a recombinant GUS baculovirus produced by using conventional recombination techniques.

10 GUS protein expressed by the recombinant GUS baculovirus of the present invention is visualized by SDS gel electrophoresis and its levels of GUS expression are compared with those of GUS expression of a recombinant GUS baculovirus produced by using conventional recombination techniques. Briefly, Sf9 insect cells are infected with the
15 recombinant GUS baculovirus of the present invention. After 2 days, the infected cells are lysed in a solution of: 20mM Tris pH8, 150 mM NaCl, 0.5% NP40 (10ml of solution per gram of cells). The lysate is loaded and run on an SDS-PAGE gel and stained with Coomassie blue. The intensity the 75Kd GUS band from the recombinant GUS baculovirus
20 produced by using the method of the present invention is compared with that from a recombinant GUS baculovirus produced by using conventional recombination techniques.

25 It will be apparent to those skilled in the art that various modifications and variations can be made in the compounds, compositions, kits, and methods of the present invention without departing from the spirit or scope of the invention. Thus, it is intended that the present invention covers the modifications and variations of this invention provided may come within the scope of the appended claims
30 and their equivalents.

CLAIMS

What is claimed is:

- 5 1. A library of double-stranded circular donor DNA, comprising:
 a donor DNA sequence which varies within a library of donor
 DNA sequences;
 a donor recombination site; and
 a selectable marker,
10 wherein the circular donor DNA does not include an origin of
 replication.
2. The library of circular donor DNA according to claim 1, wherein
 the library of donor DNA is a library of cDNA or genomic DNA.
- 15 3. The library of circular donor DNA according to claim 2, wherein
 the cDNA library is derived from single human chromosomes.
4. The library of circular donor DNA according to claim 1, wherein
20 the circular donor DNA further comprises a promoter sequence that
 controls expression of the donor DNA sequence.
5. The library of circular donor DNA according to claim 4, wherein
 the promoter is derived from bacteria, yeast, insect, animal, plant or
25 virus.
6. The library of circular donor DNA according to claim 4, wherein
 the promoter is selected from the group consisting of *E. coli lac* and *trp*
 operons, the *tac* promoter, the bacteriophage λ *p^L* promoter,
30 bacteriophage T7 and SP6 promoters, β -actin promoter, insulin
 promoter, human cytomegalovirus (CMV) promoter, HIV-LTR, RSV-LTR,

SV40 promoter, baculoviral polyhedrin and p10 promoter.

7. The library of circular donor DNA according to claim 4, wherein the promoter is an inducible promoter.

5

8. The library of circular donor DNA according to claim 7, wherein the inducible promoter is selected from the group consisting of tetracycline, heat shock, steroid hormone, heavy metal, phorbol ester, adenovirus E1A element, interferon, and serum inducible promoters.

10

9. The library of circular donor DNA according to claim 1, wherein the donor recombination site is a DNA sequence recognized by a site-specific recombinase to facilitate site-specific fusion between the circular donor DNA and an acceptor vector containing an acceptor recombination site.

15

10. The library of circular donor DNA according to claim 1, wherein the donor recombination site a recombination site recognized by a recombinase, a transposase or an integrase.

20

11. The library of circular donor DNA according to claim 1, wherein the donor recombination site is a lox site that is recognized by the Cre recombinase of bacteriophage P1.

25

12. The library of circular donor DNA according to claim 11, wherein the donor recombination site is selected from the group consisting of loxB, loxL, loxR, loxP, loxP3, loxP23, lox Δ 86, lox Δ 117, loxP511, and loxC2.

30

13. The library of circular donor DNA according to claim 11, wherein the the donor recombination site is selected from the group consisting of

att sites recognized by the Int recombinase of bacteriophage λ , a FRT site recognized by FLP recombinase of the 2pi plasmid of *Saccharomyces cerevisiae*, a recombination site recognized by the resolvase family, and a recombination site recognized by transposase of *Bacillus thuringiensis*.

14. The library of circular donor DNA according to claim 1, wherein the selectable marker is selected from the group consisting of β -galactosidase, fluorescent protein, secreted form of human placental alkaline phosphatase, β -glucuronidase, antibiotic resistance genes, yeast selectable markers leu2-d and URA3, apoptosis resistant genes, and antisense oligonucleotides.

15. The library of circular donor DNA according to claim 1, wherein the circular donor DNA further includes an affinity tag.

16. The library of circular donor DNA according to claim 15, wherein the affinity tag is selected from the group consisting of a polyhistidine tract, polyarginine, glutathione-S-transferase, maltose binding protein, a portion of staphylococcal protein A, protein A, and epitope tag.

17. The library of circular donor DNA according to claim 15, wherein the affinity tag is an EE tag.

18. The library of circular donor DNA according to claim 1, wherein the circular donor DNA further includes a polyadenylation signal.

19. A double-stranded circular donor DNA, comprising:
a donor DNA sequence;
a donor recombination site; and
at least one selectable marker, the circular donor DNA not

including an origin of replication.

20. A circular acceptor vector, comprising:

an origin of replication; and

5 an acceptor recombination site capable of recombining with a donor DNA, the acceptor vector not including a promoter for regulating expression of the donor DNA.

10 21. The circular acceptor vector according to claim 20, wherein the circular acceptor vector is capable of transforming, transfecting or transducing a host cell selected from the group consisting of a bacterial, yeast, animal, plant, and insect cell.

15 22. The circular acceptor vector according to claim 20, wherein the circular acceptor vector is a prokaryotic plasmid.

23. The circular acceptor vector according to claim 22 further comprising: a prokaryotic termination sequence selected from the group consisting of the T7 termination sequence, the bacteriophage λ
20 T_{INT} , T_{L1} , T_{L2} , T_{L3} , TR_1 , TR_2 , and T_{6S} termination signals.

24. The circular acceptor vector according to claim 19, wherein the circular acceptor vector is a mammalian expression vector.

25 25. The circular acceptor vector according to claim 24 wherein the mammalian expression vector contains one or more eukaryotic marker genes, a eukaryotic transcriptional and translational termination signal and a polyadenylation signal.

30 26. The circular acceptor vector according to claim 24 wherein the mammalian expression vector includes an origin of replication and a

selectable marker gene that functions in bacterial cells.

27. The circular acceptor vector according to claim 19 wherein the circular acceptor vector is a yeast expression vector.

5

28. The circular acceptor vector according to claim 27 wherein the yeast expression vector is selected from the group consisting of episomal vector, plasmid vector, integrating vector, and yeast chromosomes.

10

29. The circular acceptor vector according to claim 20 wherein the vector is a baculoviral vector.

15

30. The circular acceptor vector according to claim 28 wherein the baculoviral vector is a modified *Autographa californica multiple nuclear polyhedrosis virus*.

20

31. The circular acceptor vector according to claim 30 wherein the modified *Autographa californica multiple nuclear polyhedrosis virus* does not include a polyhedrin promoter.

25

32. A kit for generating a recombinant expression vector, comprising:
one or more linear donor DNA comprising a donor DNA sequence;

a linear driver DNA comprising a promoter sequence, a recombination site, and at least one selectable marker, ligation of the linear donor DNA and the linear driver DNA resulting in a circular donor DNA; and

30

a circular acceptor vector comprising an origin of replication and an acceptor recombination site capable of recombining with the circular donor DNA.

33. A kit for generating a library of recombinant expression vectors, comprising:

- 5 a library of double-stranded circular donor DNA comprising a donor DNA sequence which varies within a library of donor DNA sequences, a donor recombination site, and at least one selectable marker, the circular donor DNA not including an origin of replication; and
- 10 a circular acceptor vector comprising an origin of replication and an acceptor recombination site capable of recombining with the circular donor DNA.

34. A method for generating recombinant expression vector, comprising: contacting a circular double-stranded donor DNA and a

15 circular acceptor vector in the presence of a recombinase under conditions suitable for the circular double-stranded donor DNA and the circular acceptor vector to recombine to form a single fused circular vector,

- 20 the circular double-stranded donor DNA comprising a donor DNA sequence, a donor recombination site, and at least one selectable marker, but not including an origin of replication, and
- the circular acceptor vector comprising an origin of replication and an acceptor recombination site capable of recombining with the circular donor DNA.

25 35. The method according to claim 34, wherein the circular donor DNA further includes a promoter for regulating expression of the donor DNA.

30 36. The method according to claim 34, wherein the recombination sites on the circular donor DNA and the circular acceptor vector are both

lox sites.

37. The method according to claim 34 further comprising steps of transforming, transfecting or transducing a host cell; and
5 selecting the correctly fused recombinant vector based on the selectable phenotype conferred by the selectable marker gene on the recombinant vector.

38. The method according to claim 37 further comprising:
10 selecting the correctly fused recombinant vector by using a biotin-labeled oligonucleotide which is capable of hybridizing with circular double-stranded donor DNA.

39. The method according to claim 34, wherein the recombinase is
15 selected from the group consisting of the bacteriophage P1 Cre recombinase, yeast FLP recombinase, Inti integrase, bacteriophage λ , phi 80, P22, P2, 186, and P4 recombinase, Tn3 resolvase, the Hin recombinase, the Cin recombinase, *E. coli* xerC and xerD recombinases, *Bacillus thuringiensis* recombinase, TpnI and the β -
20 lactamase transposons, and the immunoglobulin recombinases.

40. A method for generating recombinant expression vectors, comprising: ligating one or more double-stranded linear donor DNA which includes a donor DNA sequence with a double-stranded linear
25 driver DNA which includes a promoter sequence and a donor recombination site to form a single circular donor DNA, the circular donor DNA not including an origin of replication, the donor DNA sequence being under the transcriptional control of the promoter; and
contacting the circular donor DNA and a circular acceptor vector
30 in the presence of a recombinase to form a single fused circular vector, the circular acceptor vector comprising an origin of replication and an

acceptor recombination site capable of recombining with the circular donor DNA.

5 41. The method according to claim 40, wherein the linear donor DNA and linear driver DNA contain matching restriction sites.

42. A method for generating recombinant expression vectors, comprising: ligating a library of double-stranded linear donor DNAs, where each member of the library includes a donor DNA sequence, with
10 a double-stranded linear driver DNA which includes a promoter sequence and a donor recombination site to form a single circular donor DNA, the single circular donor DNA not including an origin of replication, the donor DNA sequence being under the transcriptional control of the promoter; and

15 contacting the circular donor DNA and a circular acceptor vector in the presence of a recombinase to form a single fused circular vector, the circular acceptor vector comprising an origin of replication and an acceptor recombination site capable of recombining with the circular donor DNA.

20

43. The method according to claim 42, wherein the library of double-stranded linear donor DNAs are DNAs amplified from a library of cDNA clones.

25 44. The method according to claim 43, wherein the library of cDNA clones is arrayed in a multi-well plate.

45. The method according to claim 43, wherein the library of cDNA clones is a cosmid or phage library.

30

46. The method according to claim 42, wherein ligating the library of

double-stranded linear donor DNAs with a double-stranded linear driver DNA is performed by ligation independent cloning.

1/2




FIGURE 1

2/2




FIGURE 2

SuperGen.239.ST25

SEQUENCE LISTING

<110> Clark, Robin

<120> COMPOSITIONS AND METHODS FOR GENERATING EXPRESSION VECTORS THROUGH SITE-SPECIFIC RECOMBINATION

<130> 12636-239

<150> US 09/606,323

<151> 2000-06-28

<160> 5

<170> PatentIn version 3.0

<210> 1

<211> 34

<212> DNA

<213> Artificial sequence: loxP

<400> 1

ataacttcgt ataattgtatg ctatacgaag ttat
4

3

<210> 2

<211> 34

<212> DNA

<213> Artificial sequence: loxP511

<400> 2

ataacttcgt atagtataca ttatacgaag ttat
4

3

<210> 3

<211> 34

<212> DNA

<213> Artificial sequence: loxC2

<400> 3

acaacttcgt ataattgtatg ctatacgaag ttat
4

3

<210> 4

<211> 34

<212> DNA

<213> Artificial sequence: FLP

<400> 4

SuperGen.239.ST25

gaagttccta ttctctagaa agtataggaa cttc
4

3

<210> 5
<211> 10
<212> PRT
<213> Artificial sequence: EE-tag

<400> 5

Met Glu Glu Glu Glu Tyr Met Pro Met Glu
1 5 10

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.